

Design of the Martian far-IR ORE Spectrometer MIRORES

J. Ciężela^[1], J. Bańska^[1], J. Barylak^[1], M. Ciężela^[1], M. Kowaliński^[1], S. Płocieniak^[1], J. Gurgurewicz^[1], D. Mège^[1], B. Pieterek^[2], Ż. Szaforz^[1], P.-A. Tesson^[1], M. Giuranna^[3], F. Pirajno^[4], T. Mrozek^[1], G. Paślawski^[5], N. Zalewska^[6], M. Józefowicz^[7]

^[1]Space Research Centre, Polish Academy of Sciences, ul. Bartycka 18A, 00-716 Warsaw, Poland (jc@cbk.pan.wroc.pl)
^[2]Institute of Geology, Adam Mickiewicz University, ul. Bogumiła Krygowskiego 12, 60-680 Poznan, Poland
^[3]Institute for Space Astrophysics and Planetology (IAPS), Via del Fosso del Cavaliere 100, 00133 Roma, Italy
^[4]Centre for Exploration Targeting, The University of Western Australia, 35, Stirling Highway, Crawley WA 6009, Australia

^[5]Wroclaw University of Science and Technology, ul. Wyb. Wyspiańskiego 27, 50-370 Wroclaw, Poland
^[6]Institute of Aviation, al. Krakowska 110/114, 02-256 Warsaw, Poland
^[7]ABM Space Sp. z o.o., ul. Prosta 5/2, 87-100 Toruń, Poland




1 Introduction:

Identification of sulfide ores, being a major source of **noble metals** (Au, Ag, Pt) and **base metals** (Cu, Pb, Zn, Sn, Co, Ni, occasionally Fe), will be vital to self-sustainment of future Mars colonies. Martian meteorites are rich in sulfides^[1,2], which reflects in recent findings from Martian rovers^[3,4,5]. Yet the only high resolution (18 m/px) infrared spectrometer orbiting Mars, CRISM (Compact Reconnaissance Imaging Spectrometer for Mars) on board Mars Reconnaissance Orbiter (MRO), struggles to detect sulfides on the martian surface. Spectral interferences with silicates impede sulfide detection in the 0.4-3.9 μm CRISM range. In contrast, at least four common sulfides on Earth and Mars (pyrite, chalcopyrite, marcasite, pyrrhotite) possess prominent absorption peaks in a narrow far-infrared (FIR) wavelength range of 23-28 μm . Providing global distribution and chemical composition of sulfide ores would help to choose useful targets for future Mars exploration missions. Therefore, we are starting to design a relatively **inexpensive and simple pyroelectric detector-based Martian far-IR ORE Spectrometer (MIRORES)**.



Pyrite, Navajun, Spain; John Harrison

Sulfide spectra:



adapted from Brusentova et al., 2012

2 Design:

We propose a relatively **inexpensive, simple instrument** measuring only limited 18-40 μm range of the FIR spectrum. The **middle band** is 23-28 μm , where we expect absorption peak for ore minerals. **Additional bands**, 18-21 and 35-40 μm , will allow interpolating the background level of radiance at the middle band, which varies as a function of incidence angle, emission angle, atmospheric H₂O and other factors. **Dimensions are reduced to a size (47 x 32 x 32 cm) and mass (~10 kg) of a microsatellite.**


Detection system:

Three filters:



- (1) one for the main band of 23-28 μm
- (2,3) two for reference bands at 18-21 μm and 35-40 μm

Detector parameters:


- responsivity 1.3 V/W
- incident power limit 5 W/cm²
- operating temperature from -55 to +125°C



Three Pyroelectric Laser Detectors Model 420



Optical system:



Cassegrain optical system with large primary and small secondary mirrors

3 Summary:

The largest challenge related to this design is the small field of view conditioned by high resolution required for this study (10-20 m/px), which in limited space can be only achieved by the use of the **Cassegrain optical system**. The probe may be launched as a piggyback mission with a larger satellite between 2026 and 2030, for example **Japanese Martian Moons Exploration satellite of JAXA (2024/26)** or **Mars Sample Return Mission (HX-2) of CNSA (2028/30)**.

References

^[1] Wang, Z. & Becker, H. Chalcophile elements in Martian meteorites indicate low sulfur content in the Martian interior and a volatile element-depleted late veneer. Earth Planet. Sci. Lett. 463, 56-68 (2017)
^[2] Wang, Z. & Becker, H. Silver contents and Cu/Ag ratios in Martian meteorites and the implications for planetary differentiation. Geochim. Cosmochim. Acta 216, 96-114 (2017)
^[3] Sanyal, S. W. et al. Pyroclastic activity at home plate in Gusev crater, Mars. Sci. 316, 738-742 (2007)

^[4] Morris, R. V. et al. Iron mineralogy and aqueous alteration from Husband Hill through Home Plate at Gusev Crater, Mars: Results from the Mössbauer instrument on the Spirit Mars Exploration Rover. J. Geophys. Res. E Planets 113, E12542 (2008)
^[5] Vaniman, D. T. et al. Mineralogy of a Mudstone at Bay, Yellowknife Crater, Gale. Sci. 343, 1-9 (2014)
^[6] Brusentova et al. Laboratory far-infrared spectroscopy of terrestrial sulphides to support analysis of cosmic dust spectra. Mon. Not. R. Astron. Soc. 420, 2569-2579 (2012)

Acknowledgements

J. Ciężela is supported within the START program of the Foundation for Polish Science. D. Mège, J. Ciężela, M. Ciężela, J. Gurgurewicz and P. Tesson are supported within the EXOMHYDR project funded by the TEAM program of the Foundation for Polish Science (TEAM/16-3/20) co-financed by the European Union under the European Regional Development Fund. B. Pieterek is supported within European Funds Smart Growth (PO WER) project no. POWR.03.02.00-00-1027/17.

