ANALYSIS OF YOHKOH-OBSERVED LIMB FLARES ACCOMPANYING CME EVENTS

B. Sylwester and J. Sylwester

Space Research Centre, Polish Academy of Sciences, Wroclaw, Poland

ABSTRACT

We have made the literature/web search in order to find well observed limb flares seen by *Yohkoh* instruments. For the purpose of this study we have looked for flares associated with CME counterparts. Only few such events have been found and analysed (including compact and arcade type flares as seen on SXT images). In this contribution we present results illustrating several aspects of selected flare evolution. The results shown concern the morphology (SXT deconvolved images, HXT reconstructed images) and the flaring plasma dynamics as seen in transversal (from image cadences) and radial directions (from BCS spectral analysis). As the selected event has been observed over the limb, detailed investigation of vertical motions of the hot plasma is possible.

Key words: solar physics; flares; X-rays; CME.

1. INTRODUCTION

The *Yohkoh* collected a large number of data for flares and active regions. This offers the possibility of studying various characteristics of high temperature plasma using the four instruments placed onboard this satellite. The flaring plasma morphology can be studied in the soft and hard X-ray bands using the data from Soft X-ray Telescope (SXT) and Hard X-ray Telescope (HXT) respectively. The SXT is described in the paper by Tsuneta et al., (1991). In order to increase spatial resolution on the images, we routinely deconvolve the SXT images with oversampling. In this respect we use ANDRIL algorithm described in the papers by Sylwester and Sylwester (1998, 1999). The deconvolution removes the image blurring (due to the instrument point spread function) and reveals fine details ($\sim$1 arcsec) of flaring structure. The size of subpixel on the deconvolved image is $\sim$0.5 arcsec (oversampling factor equal 5). The HXT images are reconstructed using the maximum entropy method, MEM (Gull and Daniell, 1978; Kosugi et al., 1991). The plasma dynamics is studied based on the analysis of soft X-ray spectra obtained from the Bent Crystal Spectrometer (BCS). The spectra analysed contain the lines of Fe XXV, Ca XIX, and S XV ions observed with high spectral resolution (Culhane et al., 1991).

In this contribution we analyse a number of limb flares apparently related with coronal mass ejections (CME). Near - limb location of flares is useful for a precise coalignment of images obtained in the soft and hard X-rays. This precise coalignment is of primary importance for determining the distribution of plasma parameters using "filter ratio" technique. Standard alignment procedures adopted within the *Yohkoh* analysis software appear not adequate in this respect. As an example of the analysis performed we have selected to discuss here the event comprised of C7.0 and M2.5 importance flares observed over the limb on 16 October 2000. The first flare (precursor?) although much fainter in soft X-rays exhibits substantial nonthermal signature - a sharp, event dominating, hard X-ray spike observed up to $\sim$50 keV. The event-associated CME has been launched around the time (rise phase) of the second flare. In Figure 1 we present the three lightcurves: as seen by *GOES* ($\lambda = 1$ Å - 8 Å channel), this observed in the BCS resonance w line of Ca XIX ion and the HXT M1 channel (23 keV - 33 keV). The lightcurves displayed in the Figure embrace both flares. The first flare started at 05:32 UT and reached the maximum at 05:49 UT. The second flare began at 06:40 UT and reached the maximum at 07:28 UT. Both flares seemingly occurred at the same active region which already moved far behind the solar limb.

We have studied the evolution and morphology of brightness patterns seen on hard, MEM reconstructed HXT images, and soft, deconvolved SXT images. We have performed the analysis of time-dependent maximum brightness location for soft and hard X-rays. We also looked into the relationship between their relative positions.

To supplement the analysis of transversal variations (which come as a result of image analysis), we have used the BCS spectra obtained in the vicinity of resonance lines of Fe XXV, Ca XIX and S XV ions, for determination of the radial motion in the observed flares. In addition, observed changing line broadening contains information on nonthermal (turbulent) plasma motions. By fitting a Gaussian profile to the



Figure 1. The lightcurves for the double event on 16 October 2000. The plots represent: 1 Å - 8 Å emission seen by GOES *(smooth black line over entire time interval), Ca* XIX *resonance line emission as observed by BCS (separate grey curves covering four* Yohkoh *orbits) and M1 channel emission of HXT (23 keV - 33 keV) for two orbits when this emission was above background. Note the reverse proportions between soft and hard X-ray emission among the two flares. The curves plotted are normalised (in respect to their maxima) in order to bring them to common scale.*

resonance lines of Fe XXV, Ca XIX and S XV ions we have determined respective line positions and line widths. For more comprehensive description of the analysis of directed and turbulent plasma motions the reader is referred to papers by Fludra et al., (1989), Doschek et al., (1993) and Sterling et al., (1996).

As is seen in Figure 1 the BCS data are available for analysed flares also towards the late decay phase. The spectral line positions as determined for this late decay phase have been treated by us as reference, i.e. we assumed no directed plasma motions at that time. The results presented do not depend heavily on this assumption. As the SXT and BCS images show-up presence of a hotter ($T > 5$ MK) thermal plasma (Sylwester and Sylwester, 2002) and the emission seen in higher channels on HXT *Yohkoh* images may come also from the nonthermal contributors, using these data we can look into the thermal/nonthermal plasma relationship.

2. THE FIRST FLARE ON 16 OCTOBER 2000

Inspection of sequences of deconvolved SXT images shows that the overall shape of the soft X-ray emission seen in AlMg, Al12 and Be119 channels share similar patterns as concerns both spatial and time variations. The emission is at the beginning concentrated in chains of small localised kernels "protruding" just above the limb into the corona. This chains resemble the loop-like structure as seen in lower resolution. Later on, the brightest emission moves a bit higher being concentrated at one time (05:41 UT) in the northern leg of the loop. During the soft X-ray maximum (towards the end of hard X-ray phase) the most prominent becomes compact kernel located at



Figure 2. The composite picture showing SXT Be119 deconvolved images (top row), coinciding MEM reconstructed HXT M1 (23 keV - 33 keV) images (middle row) for three peaks of hard X-ray emission. This three peaks make input to the first spike seen in Figure 1. The soft (GOES 0.5 Å - 4 Å) and hard (HXT LO and M1) X-ray lightcurves are displayed in the bottom row for the comparison. The hatched areas mark time intervals used for HXT image reconstruction.

(what appears to be) the southern leg's base. Some of this changing pattern may be seen in the upper panel of Figure 2. Note that the intensities on all images have been normalised to the local maximum on each image. During the hard X-ray rise phase the bright soft X-ray patches tend to "avoid" regions of increased nonthermal hard X-ray emissions. This is best seen in the initial frames (first column presented in Figure 2) where the soft X-ray emission encompasses the hard X-ray emission kernel. After the maximum of the hard X-ray emission, the soft and hard X-ray emission patches appear to be nearly cospatial in projection which is seen in Figure 2 (last column) and in detail in Figure 3. There, the exact overlay of three emission patterns simultaneously obtained during the hard X-ray post-maximum phase is presented. The images in Be119 and in HXT bands have been summed over the same time interval in order to improve the statistic. The cospatiality of the emission seen at this time rules out any possibility that the differences in location of the soft and hard X-ray sources seen before were related to uncertainties of the coalignment. The SXT and HXT images have been coaligned using *Yohkoh* pointing data files and the *SolarSoft* routines. The coalignment accuracy of SXT and HXT images is therefore anticipated to be of the order of ~ 1 arcsec. From the inspection of Figure 3 and the other available SXT and HXT pairs of thus overlaid images, it appears that during the rise phase of hard X-ray emission the hard and



Figure 3. *The overlay of three emission patterns observed simultaneously at the phase just after the main hard X-ray maximum (at ∼05:44:08 UT). The deconvolved SXT Be119 emission is placed as background with the white thin contours drawn at the indicated levels. The HXT LO-band emission (14 keV - 23 keV) is represented by black dashed contours and HXT M1 emission, (23 keV - 33 keV) by solid black contour.*



Figure 4. *The evolution of maximum brightness location (contours drawn at 0.85 of maximum intensity) for consecutive frames as seen on deconvolved Be119 images. The contours are overlaid atop the background emission pattern (in grey) representing the first Be119 image (05:35:25 UT) obtained in the flare SXT mode. The vertical bar to the right of the image gives the grey shade - time relation.*

soft X-ray emission comes from separate plasma locations (volumes). This is rather surprising since for most of flares analyzed yet using the same technique, the HXT and SXT emission patches although not exactly cospatial, have not been separated so much as in the present case. During the hard X-ray decay phase of this flare the soft and hard emissions come from regions overlapping in projection. From the discussed picture showing different location of the harder and softer X-ray sources, the physical links between them are not obvious.

It is convenient to study the soft X-ray bright kernel's location by looking into time sequence of contours drawn at the level close to the maximum (0.85 of maximum intensity in the present case). From such a sequence plot shown in Figure 4 it is seen, that the most intense soft X-ray emission changes its position considerably. The individual bright kernels become visible at different locations during ∼15 minutes of evolution (traced by SXT Be119 flare sequence). The displacements observed are as much as ∼40 arcsec in the direction along the solar limb and ∼25 arcsec radially. The contours become lighter as time passes towards the end of ∼15 minutes time interval considered. It is seen that the emission observed towards the end is placed very close (loop top?) to a pair of emission kernels seen at the beginning. Meantime however, the brightest kernels are located quite far away from that place (forming quasi loop-like structure 20 000 km high). The colour animation showing the evolution pattern presented in Figure 4 can be downloaded from: `www.cbk.pan.wroc.pl/publications/2003/cme_accomp_flare/be_cont_085_A.avi`.



Figure 5. *Results of the analysis of BCS spectra as obtained for the first flare. In the top panel the time variations of GOES X-ray flux in 1 Å - 8 Å band is shown as smooth, flat curve. The HXT M1 emissions in 23 keV - 33 keV is over plotted (spiky line). Both curves are normalised over the period covering both flares and therefore the GOES level is relatively low. In the middle panel the positions of resonance line centroids are plotted for Fe XXV (asterisks), Ca XIX (triangles) and S XV ions (circles). At the bottom panel the variations of respective lines' FWHMs are presented.*

Using the spectra observed by the BCS we have studied the flaring plasma dynamics. In this respect we looked into variation of the centroid of line position and the line width. We have used the resonance lines of Fe XXV, Ca XIX and S XV ions and performed the